
NROSCI/BIOSC 1070 and MSNBIO 2070 Exam # 1 September 27, 2019

Each letter can


be

(2



Ventricular blood volume constitutes preload: f)


- 2) Following a heart attack, a patient is suffering from heart failure.
 - a) On the diagram below, indicate how the Starling curve changes from normal on the side of the heart that is damaged. *(3 points)*.
 - **b)** The drug digoxin is sometimes prescribed to treat heart failure. The primary mechanism of action involves inhibition of Na+/K+ ATPase, mainly in the myocardium. On the diagram below, indicate how digoxin alters the Starling curve for the damaged ventricle. *(3 points).*



c) Briefly describe the physiologic mechanism through which digoxin causes the change in the Starling curve you indicated above. *(6 points).*

- **3)** An individual is administered phenylephrine, an α -1 receptor agonist. Assume that there are no compensatory or reflex-elicited changes that alter the direct effects of the drug on the cardiovascular system.
 - a) Would the α -1 receptor agonist result in a change in blood pressure? If so, would blood pressure increase or decrease? What physiological actions of phenylephrine cause this change in blood pressure? (4 points).

b) Normal vascular and cardiac function curves are illustrated below. Indicate how the administration of an α -1 receptor agonist would alter the curves. You may also add a description to clarify your response. *(6 points).*



4) Does the sympathetic nervous system have any effect on skeletal muscle? If so, describe the receptors on skeletal muscle through which the sympathetic nervous system elicits actions, and the physiologic effects of the sympathetic nervous system on skeletal muscle. *(5 points).*

5) It is not uncommon for cranial nerve III (the oculomotor nerve) to be damaged by an aneurysm from a large artery in the head, the posterior communicating artery. Often, the parasympathetic fibers in the nerve are damaged early. What are the consequences of damage to the parasympathetic nerve fibers in the third cranial nerve? (6 points).

- 6) Binding of an agonist to either GABAa or glycine receptors increases the conductance of the same ion through the membrane.
 - a) Which ion is transported more readily through the membrane following the binding of agonists to these receptors? *(3 points).*

b) What is the physiologic effect of an agonist binding to either GABAa or glycine recptors? (What is the effect on membrane potential?) *(3 points)*.

c) Are GABAa and glycine receptors characterized as metabotropic or ionotropic receptors (circle the answer below)? *(2 points).*

Metabotropic

Ionotropic

d) Are the synaptic effects of GABA and glycine terminated by breaking down the transmitters or reuptaking them into nerve terminal? (2 points).

Reuptake

Breakdown

- 7) The following questions relate to muscle unit types. Circle which muscle unit type best meets the stated criterion. *(1 point each; 6 points total).*
 - Most ATP usage per unit time a) FF S Most actin and myosin content per muscle cell b) FF S C) Most similar to cardiac muscle cells FF S Most able to undergo substantial hypertrophy (increase in diameter by d) adding actin and myosin) FF S e) Produce the most tension during contraction FF S Can contract for a sustained period without fatigue f) FF S

8) A critically ill patient is given dobutamine at a dose that mainly serves as a β -1 receptor agonist. What effects would the drug have on the following? (circle the correct answer) (2 points each; 10 points total).

a)	End systolic volume		
	Unchanged	Higher	Lower
b)	End diastolic volume		
	Unchanged	Higher	Lower
C)	Ventricular filling time		
	Unchanged	Higher	Lower
d)	Blood pressure		
	Unchanged	Higher	Lower
e)	Workload of the heart		
	Unchanged	Higher	Lower

9) Patient A is given a selective β -1 receptor agonist while Patient B is given a drug that activates both β -1 and β -2 receptors. If both drugs have equivalent effects on β -1 receptors, which will produce the greatest change in afterload? Discuss the physiologic mechanism accounting for your answer. *(7 points).*

10) During a spinal surgery, the rostral portion of the sympathetic chain on one side is destroyed. As a result, the patient loses all sympathetic innervation of the head on that side. List three distinct physiologic changes that would result from removal of sympathetic innervation from half of the face. *(7 points).*

11) The following physiologic parameters are determined for an individual:

Systolic aortic Pressure = 150 mm Hg Diastolic aortic pressure = 90 mm Hg Systolic pulmonary artery pressure = 12 mm Hg Diastolic pulmonary artery pressure = 6 mm Hg Heart rate = 50 beats/min Left atrial pressure = 5 mm Hg Right atrial pressure = 2 mm Hg End systolic volume (both ventricles) = 50 ml End diastolic volume (both ventricles) = 150 ml

For this individual, determine the resistance in the pulmonary circulation relative to the systemic circulation (i.e., Pulmonary resistance/Systemic resistance). You MUST show your calculations to receive credit. *(10 points).*

12) A patient has highly elevated plasma levels of ACTH, β -LPH, and β -endorphin. What is the most likely cause of these abnormal plasma hormonal levels? (5 *points).*

Please Print Your Name Below This page will be filed away until exam grading is complete.

Print Your Name

Exam # 1 September 27, 2019